

This article was downloaded by: [Gian Italo Bischi]

On: 10 August 2015, At: 07:06

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: 5 Howick Place, London, SW1P 1WG



Journal of Difference Equations and Applications

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gdea20>

On a discrete-time model with replicator dynamics in renewable resource exploitation

Gian Italo Bischi^a, Lorenzo Cerboni Baiardi^b & Davide Radi^c

^a Department of Economics, Society, Politics, University of Urbino, Italy

^b Department of Base Sciences and Fundamentals, University of Urbino, Italy

^c Department of Management, Marche Polytechnic University, Ancona, Italy

Published online: 30 Jul 2015.



CrossMark

[Click for updates](#)

To cite this article: Gian Italo Bischi, Lorenzo Cerboni Baiardi & Davide Radi (2015): On a discrete-time model with replicator dynamics in renewable resource exploitation, Journal of Difference Equations and Applications, DOI: [10.1080/10236198.2015.1059830](https://doi.org/10.1080/10236198.2015.1059830)

To link to this article: <http://dx.doi.org/10.1080/10236198.2015.1059830>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

On a discrete-time model with replicator dynamics in renewable resource exploitation

Gian Italo Bischi^{a,1}, Lorenzo Cerboni Baiardi^{b*} and Davide Radi^{c,2}

^aDepartment of Economics, Society, Politics, University of Urbino, Italy; ^bDepartment of Base Sciences and Fundamentals, University of Urbino, Italy; ^cDepartment of Management, Marche Polytechnic University, Ancona, Italy

(Received 28 February 2015; accepted 29 May 2015)

We consider a discrete-time version of the model proposed by Lamantia and Radi [15] to describe a fishery where a population regulated by a logistic growth function is exploited by a pool of agents that can choose, at each time period, between two different harvesting strategies according to a profit-driven evolutionary selection rule. The resulting discrete dynamical system, represented by a two-dimensional nonlinear map, is characterized by the presence of invariant lines on which the dynamics are governed by one-dimensional restrictions that represent pure, i.e. adopted by all players, strategies. However, interesting dynamics related to interior attractors, where players playing both strategies coexist, are evidenced by analytical as well as numerical methods that reveal local and global bifurcations.

Keywords: discrete-time population model; replicator dynamics; resource exploitation; attractors; bifurcations

1. Introduction

In this paper, we consider a discrete-time version of the model proposed in [15] to describe a fishery where two different harvesting strategies can be employed, one denoted as standard and the other one more ecological (less intensive, hence more environmentally friendly). At any time period, the fish population is assumed to reproduce according to a discrete-time logistic growth function (see e.g. [7] Chapter 2 and [13]), and the agents that exploit the fishery are assumed to update their harvesting strategy according to a profit-driven adaptive mechanism based on the evolutionary selection rule known as replicator dynamics (see e.g. [14,20]).

Even if dynamic models in ecology have been traditionally formulated in continuous time, discrete-time population models have received a great amount of attention not only for the complex and intriguing dynamics that they can produce even in the simplest systems, but also because biological motivations have been proposed to explain their usefulness in ecologic modelling. In fact, several authors stress that discrete-time population models should be used whenever reproduction happens at given breeding seasons, as several animal species successfully mate only during certain times of the year, thus giving non-overlapping generations (see e.g. [12,17]). So, more and more discrete-time population models have been proposed in the literature (see e.g. [5,10,13]). Moreover, as already stressed in [15], in the model considered in this paper decisions about the kind of harvesting strategy to be adopted typically occur in discrete time, as such

*Corresponding author. Email: l.cerbonibaiardi@campus.uniurb.it

decisions imply the adoption of different fishing technologies, and/or different numbers of workers with different kinds of equipment, hence they cannot be revised at any time. Sometimes the possibility of switching from a harvesting strategy to another one is allowed only at given time periods by laws that regulate harvesting activities (see [3,4,6]).

On the basis of these motivations, a discrete-time model, represented by a two-dimensional nonlinear map, is studied in this paper by using analytical, geometric and numerical methods. The structure of the map is quite interesting from the point of view of its mathematical properties, and is typical (hence representative) for a large class of repeated evolutionary games where a population of N players can choose, at any time, between two strategies. In fact, one dynamic variable, denoted as $r(t) \in [0, 1]$, represents the fraction of players adopting a given strategy at time period $t \in \mathbb{N}$ (of course the complementary fraction $1 - r(t)$ adopts the other strategy at the same time period). As typically occurs in these kind of evolutionary games, the two lines $r = 0$ and $r = 1$, where all players adopt the same strategy, are invariant lines, along which the dynamics characterized by unique kind of players (pure strategy case) are governed by a one-dimensional restriction of the map. In our case, the dynamics along such invariant boundary lines are given by the iteration of a quadratic map, topologically conjugate to the standard logistic map. However, interior attractors, where players carrying out both strategies coexist, can be obtained, and some bifurcations involving interior and boundary invariant sets can be studied. Indeed, very rich dynamic scenarios can be highlighted, both analytically and numerically, and regions of the phase space of the model can be detected in which quasi-periodic motions prevail (i.e. where the linear approximation of the map has complex eigenvalues) and other regions where real eigenvalues give rise to monotonic motions or improper oscillations. In both cases, however, transitions to chaotic behaviours can be observed. Moreover, the existence of non-topological Milnor attractors embedded in the invariant lines is proved and numerically shown.

The paper is organized as follows. Section 2 presents the discrete-time model and discusses the dynamics along the invariant lines where one-dimensional dynamics occurs when all agents adopt the same strategy. Section 3 contains analytical results on the existence of equilibrium points and their local stability properties as well as local bifurcations. Section 4 gives some propositions on global behaviour and some numerical simulations of the model. Section 5 concludes and indicates further research issues.

2. The model

Following the general setup of the model proposed by [15], let $x(t)$ denote the available quantity at time t of a renewable resource and let us consider a population of N agents that can exploit the resource by two different technologies: a standard (intensive) one characterized by technology coefficient $q_1 > 0$, and a more environmentally friendly (let's say ecological) technology characterized by $q_0 \in (0, q_1)$. Let $r(t) \in [0, 1]$ be the fraction of agents that adopt the standard technology during time period t and consequently the complementary fraction of agents $(1 - r(t))$ adopts the ecological technology, so that $r = 0$ means that all the agents adopt the ecological technology q_0 , and $r = 1$ means that all the agents adopt the standard technology q_1 . If h_i denotes the harvesting of resource by using technology i , $i = 0, 1$, following again [15] we assume that the cost functions are given by

$$C_i(h_i) = c_i + \gamma \frac{h_i^2}{q_i x}; \quad i = 0, 1, \quad (1)$$

where $c_i \leq 0$ represents fixed costs and $\gamma > 0$ is a cost coefficient.

If we denote by $a_0 > 0$ the constant price at which consumers buy the resource harvested by ecological technology and $a_1 \in (0, a_0)$ the price at which they buy the standard one, then, as shown in [15], the optimal harvesting, computed as Nash equilibrium, of the representative player that uses technology i is given by

$$h_i(x) = \frac{a_i q_i}{2\gamma} x; \quad i = 0, 1 \tag{2}$$

In the following $a_0 > a_1$ will be assumed, i.e. the loss in efficiency of the more ecological harvesting strategy is counterbalanced by a higher price that consumers wish to pay for the more environmentally friendly product.

Differently from [15], we consider a discrete-time model, i.e. a resource is given by a population with non-overlapping generations growing according to a discrete-time logistic equation

$$x(t + 1) = x(t) + \alpha x(t) \left(1 - \frac{x(t)}{k} \right) - Nr(t)h_1(t) - N(1 - r(t))h_0(t). \tag{3}$$

Here $\alpha > 0$ is the natural growth rate of the resource, $k > 0$ represents the carrying capacity, i.e. the equilibrium level of the resource in the absence of harvesting, and fraction $r(t)$ is assumed to evolve according to an exponential replicator dynamics (see e.g. [9] and [14]) driven by profits

$$\pi_i(x) = a_i h_i(x) - c_i - \gamma \frac{h_i^2(x)}{q_i x} \tag{4}$$

and expressed by

$$r(t + 1) = \frac{r(t)e^{\beta\pi_1(t)}}{r(t)e^{\beta\pi_1(t)} + (1 - r(t))e^{\beta\pi_0(t)}} = \frac{r(t)}{r(t) + (1 - r(t))e^{\beta(\Delta\pi(t))}}, \tag{5}$$

where $\beta \in [0, +\infty)$ is the so-called intensity of choice parameter and measures the reactivity of agents to adopt the more profitable strategy and $\Delta\pi(t)$ is the difference between the two profits. According to Equations (4) and (2), we have that

$$\Delta\pi(t) = \pi_0(t) - \pi_1(t) = \frac{a_0^2 q_0 - a_1^2 q_1}{4\gamma} x(t) - \xi \tag{6}$$

with $\xi = c_0 - c_1$. The parameter $\xi \in \mathbb{R}$ represents the difference between fixed costs associated with the two technologies, and may be considered as a policy parameter as it includes taxes imposed in order to obtain the prevalence of one technology over the other. In the following, we shall mainly consider $\xi \leq 0$, assuming that fixed costs for the more intensive harvesting method are higher, due to more sophisticated technology and higher taxes, or equivalently to government subsidies for agents adopting the more ecological fishing methods.

All in all, the dynamic model can be written in the form of an iterated map of the plane $T : (x, y) \rightarrow (x', y')$ with

$$T : \begin{cases} x' = \left(1 + \alpha - \frac{Na_0q_0}{2\gamma} \right) x - \frac{\alpha}{k} x^2 + \frac{N}{2\gamma} (a_0q_0 - a_1q_1) x r \\ r' = \frac{r}{r + (1-r)e^{\beta \left(\frac{a_0^2 q_0 - a_1^2 q_1}{4\gamma} x - \xi \right)}} \end{cases}, \tag{7}$$

where $'$ denotes the unit-time advancement operator, and the dynamic variables represent feasible states of the system if $x \geq 0$ and $0 \leq r \leq 1$.

It is worth underlining that the line of resource extinction $x = 0$, as well as the two lines of pure strategies $r = 0$ and $r = 1$, are invariant sets. The dynamics along the invariant line $x = 0$ is governed by the one-dimensional restriction

$$x = 0 : \quad r' = g(r) = \frac{r}{r + (1 - r)e^{-\xi}} \quad (8)$$

which is a convex function in the interval $r \in [0, 1]$ with fixed points in $r = 0$ (stable) and $r = 1$ (unstable). The dynamics along the invariant line $r = 0$, where all the agents adopt the ecological strategy, are governed by the one-dimensional restriction

$$r = 0 : \quad x' = f_0(x) = \left(1 + \alpha - \frac{Na_0q_0}{2\gamma}\right)x - \frac{\alpha}{k}x^2 \quad (9)$$

topologically conjugate to the standard logistic map $z' = \mu z(1 - z)$ by the transformation $z = (2\gamma\alpha/(k[2\gamma(1 + \alpha) - Na_0q_0]))x$ and parameter $\mu = 1 + \alpha - Na_0q_0/(2\gamma)$. Its two fixed points are given by $x_0^0 = 0$ (extinction equilibrium) and

$$x_0^* = \frac{k(2\alpha\gamma - Na_0q_0)}{2\alpha\gamma} \quad (10)$$

that represents the viable equilibrium under ecological harvesting. Notice that x_0^* is stable for the dynamics along the line $r = 0$ provided that

$$\alpha - 2 < \frac{Na_0q_0}{2\gamma} < \alpha, \quad (11)$$

where the condition $Na_0q_0/(2\gamma) = \alpha$ represents the transcritical bifurcation along the line $r = 0$ at which the viable equilibrium x_0^* merges with the extinction equilibrium x_0^0 , whereas the condition $Na_0q_0/(2\gamma) = \alpha - 2$ represents a period-doubling bifurcation, at which a stable cycle becomes the unique attractor along the line $r = 0$. As it is well known, this bifurcation opens the period-doubling cascade, leading to chaotic motion along the line $r = 0$, as the aggregate parameter $Na_0q_0/(2\gamma)$ is further decreased.

Analogously, along the invariant line $r = 1$ where all agents adopt the standard (more intensive) fishing strategy, the dynamics are governed by the map

$$r = 1 : \quad x' = f_1(x) = \left(1 + \alpha - \frac{Na_1q_1}{2\gamma}\right)x - \frac{\alpha}{k}x^2 \quad (12)$$

conjugate to the logistic map $z' = \mu z(1 - z)$ by the transformation $z = (2\gamma\alpha/(k[2\gamma(1 + \alpha) - Na_1q_1]))x$ and parameter $\mu = 1 + \alpha - Na_1q_1/(2\gamma)$. Here, the viable equilibrium is

$$x_1^* = \frac{k(2\alpha\gamma - Na_1q_1)}{2\alpha\gamma}. \quad (13)$$

Notice $x_0^* > x_1^*$ if $a_0q_0 < a_1q_1$, a condition that we shall assume in the following in order to characterize the technology q_0 as more ecological. As for x_0^* , stability conditions of the viable equilibrium along the line $r = 1$ can be obtained by straightforward

calculations, and are given by

$$\alpha - 2 < \frac{Na_1q_1}{2\gamma} < \alpha, \tag{14}$$

where similar statements about the transcritical and period-doubling bifurcations hold.

The existence of these invariant lines that bound the two-dimensional phase space of the dynamical system Equation (7) is important in order to characterize its global dynamical properties. Moreover, the knowledge of the kind of dynamic motion occurring along the two lines where a single pure strategy exists, tell us what will happen in the long run when one of the two strategies becomes dominant in terms of profits so that it will prevail due to evolutionary pressure. The latter problem may be equivalently stated by asking when the one-dimensional attractors of the restrictions along the invariant lines $r = 0$ and $r = 1$ given by Equations (9) and (12) respectively, are also attractors of the two-dimensional dynamical system. This depends on the transverse stability as well as on the existence of attractors interior to the phase space, i.e. characterized by $r \in (0, 1)$. These are the questions examined, analytically and numerically, in the next sections. Here, for the sake of clarity, it is worth specifying that the j -cycle ($j \geq 1$) laying on an invariant line has one of its eigenvectors that is along the invariant line itself, while the other eigenvector has generally another direction. This last eigenvector is commonly named transverse eigenvector, which is tangent to the so-called transverse invariant manifold.

3. Existence and stability of equilibrium points

The equilibrium points of the model Equation (7) are solutions of the system

$$\begin{cases} x \left[\alpha - \frac{Na_0q_0}{2\gamma} - \frac{\alpha}{k}x + \frac{N}{2\gamma}(a_0q_0 - a_1q_1)r \right] = 0 \\ r(1-r) \left[e^{\beta \left(\frac{a_0^2q_0 - a_1^2q_1}{4\gamma}x - \xi \right)} - 1 \right] = 0 \end{cases} \tag{15}$$

The extinction fixed points $E_0^0 = (0, 0)$ and $E_1^0 = (0, 1)$ always exist. Moreover, if $\xi = 0$ then any point of the whole segment $(0, r)$, with $r \in [0, 1]$, is a fixed point. Other boundary equilibrium points are $E_0^* = (x_0^*, 0)$ with x_0^* given by Equation (10) and $E_1^* = (x_1^*, 1)$ with x_1^* given by Equation (13). Furthermore, an interior equilibrium may exist, characterized by the co-existence of both harvesting strategies, given by $E^* = (x^*, r^*)$ with

$$x^* = \frac{4\gamma\xi}{a_0^2q_0 - a_1^2q_1}; \quad r^* = \frac{2\alpha\gamma(k - x^*) - Nka_0q_0}{Nk(a_1q_1 - a_0q_0)} \tag{16}$$

provided that $x^* > 0$ and $r^* \in (0, 1)$.

Let us notice that if $\xi < 0$, i.e. $c_0 < c_1$ as argued above, then $x^* > 0$ provided that $a_0^2q_0 < a_1^2q_1$, which is a more stringent condition than $a_0q_0 < a_1q_1$ being $a_0 > a_1$. We shall assume that this condition is satisfied in the following³. It is worth noticing that the condition $r^* \in (0, 1)$ can be easily expressed in term of the carrying capacity k , as

$r^* = 0$ for $k = k_0$ with

$$k_0 = \frac{2\alpha\gamma x^*}{2\alpha\gamma - Na_0q_0} = \frac{8\alpha\gamma^2\xi}{(2\alpha\gamma - Na_0q_0)(a_0^2q_0 - a_1^2q_1)} \quad (17)$$

and $r^* = 1$ for $k = k_1$ with

$$k_1 = \frac{2\alpha\gamma x^*}{2\alpha\gamma - Na_1q_1} = \frac{8\alpha\gamma^2\xi}{(2\alpha\gamma - Na_1q_1)(a_0^2q_0 - a_1^2q_1)} \quad (18)$$

with $k_0 < k_1$ being $a_0q_0 < a_1q_1$, so that $r^* \in (0, 1)$ for $k_0 < k < k_1$.

These existence conditions are the same as the ones given in [15] for the model in continuous time, whereas the stability conditions now are different. In order to study the local stability of the equilibrium points, we consider the Jacobian matrix

$$J(x, r) = \begin{bmatrix} 1 + \alpha - \frac{Na_0}{2\gamma} - \frac{2\alpha}{k}x - \frac{N}{2\gamma}(a_1q_1 - a_0q_0)r & -\frac{N(a_1q_1 - a_0q_0)x}{2\gamma} \\ \frac{\beta r(1-r)(a_1^2q_1 - a_0^2q_0)e^{\beta\Delta\pi}}{4\gamma(r+(1-r)e^{\beta\Delta\pi})^2} & \frac{e^{\beta\Delta\pi}}{(r+(1-r)e^{\beta\Delta\pi})^2} \end{bmatrix}. \quad (19)$$

At E_0^0 we have

$$J(0, 0) = \begin{bmatrix} 1 + \alpha - \frac{Na_0q_0}{2\gamma} & 0 \\ 0 & e^{\beta\xi} \end{bmatrix} \quad (20)$$

so E_0^0 is stable along the vertical direction (r direction) and stable along the horizontal one (x direction) provided that $Na_0q_0/(2\gamma) - 2 < \alpha < Na_0q_0/(2\gamma)$. At E_1^0 we have

$$J(0, 1) = \begin{bmatrix} 1 + \alpha - \frac{Na_1q_1}{2\gamma} & 0 \\ 0 & e^{-\beta\xi} \end{bmatrix} \quad (21)$$

so E_1^0 is unstable along the vertical direction and stable along the horizontal one provided that $Na_1q_1/(2\gamma) - 2 < \alpha < Na_1q_1/(2\gamma)$. At E_0^* , we have

$$J(x_0^*, 0) = \begin{bmatrix} 1 - \alpha + \frac{Na_0q_0}{2\gamma} & -\frac{N(a_1q_1 - a_0q_0)(2\alpha\gamma - Na_0q_0)}{2\alpha\gamma^2} \\ 0 & e^{\beta\left(\frac{k(a_1^2q_1 - a_0^2q_0)(2\alpha\gamma - Na_0q_0)}{8\alpha\gamma^2} + \xi\right)} \end{bmatrix} \quad (22)$$

so E_0^* is stable along the eigendirection transverse to $r = 0$ if $k < k_0$, with k_0 given by (17), and stable along the horizontal direction if $Na_0q_0/(2\gamma) < \alpha < Na_0q_0/(2\gamma) + 2$.

At E_1^* we have

$$J(x_1^*, 1) = \begin{bmatrix} 1 - \alpha + \frac{Na_1q_1}{2\gamma} & -\frac{N(a_1q_1 - a_0q_0)(2\alpha\gamma - Na_1q_1)}{2\alpha\gamma^2} \\ 0 & e^{-\beta\left(\frac{k(a_1^2q_1 - a_0^2q_0)(2\alpha\gamma - Na_1q_1)}{8\alpha\gamma^2} + \xi\right)} \end{bmatrix} \quad (23)$$

so E_1^* is stable along the eigendirection transverse to $r = 1$ if $k > k_1$ with k_1 given by Equation (18), and stable along the horizontal direction if $Na_1q_1/(2\gamma) < \alpha < Na_1q_1/(2\gamma) + 2$.

Notice that all the stability conditions along the horizontal invariant lines $r = 0$ and $r = 1$, on which the boundary-fixed points are located, correspond to those already examined for the logistic restrictions [Equations \(9\) and \(12\)](#). Finally, at E^* we have

$$J(x^*, r^*) = \begin{bmatrix} 1 - \frac{\alpha x^*}{k} & -\frac{N(a_1 q_1 - a_0 q_0)x^*}{2\gamma} \\ \frac{\beta}{4\gamma} r^*(1 - r^*)(a_1^2 q_1 - a_0^2 q_0) & 1 \end{bmatrix}. \tag{24}$$

Hence, given that trace and determinant of the matrix [Equation \(24\)](#) are respectively

$$Tr = 2 - \frac{\alpha x^*}{k} = 2 - \frac{4\alpha\gamma\xi}{k(a_0^2 q_0 - a_1^2 q_1)} \tag{25}$$

$$Det = 1 - \frac{\alpha x^*}{k} + \frac{N\beta}{8\gamma^2} (a_1 q_1 - a_0 q_0)(a_1^2 q_1 - a_0^2 q_0)x^* r^*(1 - r^*) \tag{26}$$

a sufficient condition for the local asymptotic stability of E^* is that the eigenvalues of [Equation \(24\)](#), which are solutions of the characteristic equation $P(z) = z^2 - Tr \cdot z + Det = 0$, are located inside the unit circle of the complex plane. A necessary and sufficient condition for this is given by the following system of inequalities (known as Schur or Jury's conditions, see e.g. [\[11\]](#))

$$P(1) = 1 - Tr + Det > 0; \quad P(-1) = 1 + Tr + Det > 0; \quad 1 - Det > 0. \tag{27}$$

In our case, we have $P(1) = -4N\beta\xi/(8\gamma)(a_1 q_1 - a_0 q_0)r^*(1 - r^*) > 0$ which vanishes (and then changes sign) when r^* crosses the value $r^* = 0$ from above and when r^* crosses the value $r^* = 1$ from below. These two conditions correspond to transcritical bifurcations when the interior equilibrium $E^* = (x^*, r^*)$ merges with the boundary points E_0^* and E_1^* respectively. In fact, the condition $r^*(x^*) = 0$ implies $x^* = x_0^*$ and $r^*(x^*) = 1$ implies $x^* = x_1^*$. The two bifurcation conditions can easily be expressed in terms of the carrying capacity k as $k = k_0$ and $k = k_1$, respectively, see also [Figure 1](#),




Figure 1. Fixed points $E^*(k)$, $E_0^*(k)$ and $E_1^*(k)$ are shown as $k \in [k_0, k_1]$. The point E^* merges with E_0^* in $r = 0$ for $k = k_0$ and merges with E_1^* in $r = 1$ for $k = k_1$. Other fixed points do not depend on k .

where the equilibrium points are represented for k in the range $[k_0, k_1]$. Notice that a change of sign of $P(1)$ also occurs when ξ changes from negative to positive, and this is a transcritical bifurcation as well, but of codimension two as it occurs when the fixed point E^* crosses at $\xi = 0$ the segment of fixed points along the axis $x = 0$.

Before analysing the other two stability conditions, let us consider the condition $Tr^2 - 4Det > 0$ leading to real eigenvalues. This condition becomes

$$2\alpha^2\gamma^2x^* + 16k\alpha\gamma^2 > k^2N\beta(a_1q_1 - a_0q_0)(a_1^2q_1 - a_0^2q_0)r^*(1 - r^*) \quad (28)$$

from which it is evident that it is surely satisfied (hence we have real eigenvalues) for r^* very close to 0 or r^* very close to 1, whereas it is surely not satisfied (hence we have complex conjugate eigenvalues) for intermediate values of r^* and sufficiently high values of the parameter β . Notice that both x^* and r^* do not depend on the parameter β .

Indeed, if the other parameters are fixed so that $x^* > 0$ and r^* has intermediate values, i.e. it is not too close to the invariant lines of pure strategies, then a Neimark–Sacker bifurcation occurs for increasing values of β . In fact, the third stability condition $1 - Det > 0$, where a change of sign of the left hand side indicates the occurrence of a Neimark–Sacker bifurcation of E^* , becomes

$$N\beta(a_1q_1 - a_0q_0)(a_1^2q_1 - a_0^2q_0)r^*(1 - r^*) < \frac{8\alpha\gamma^2}{k}. \quad (29)$$

This stability condition can be equivalently written as $\beta < \beta_{NS}$, with

$$\beta_{NS} = \frac{8\alpha\gamma^2}{Nk(a_1q_1 - a_0q_0)(a_1^2q_1 - a_0^2q_0)r^*(1 - r^*)} \quad (30)$$

and r^* given by Equation (16), and a Neimark–Sacker bifurcation occurs when β increases across β_{NS} .

Finally, the stability condition $P(-1) > 0$ becomes

$$32k\gamma^2 + Nk\beta(a_1q_1 - a_0q_0)(a_1^2q_1 - a_0^2q_0)x^*r^*(1 - r^*) > 16\alpha\gamma^2x^* \quad (31)$$

that can be equivalently written as $\beta > \beta_F$ with

$$\beta_F = 2\beta_{NS} - \frac{8\gamma}{N\xi(a_0q_0 - a_1q_1)r^*(1 - r^*)}. \quad (32)$$

These results can be summarized by the following statement:

Proposition 1. The map Equation (7), with $\xi \in (-\infty, +\infty)$ and positive values of all other parameters, always has the boundary fixed points $E_0^0 = (0, 0)$ and $E_1^0 = (0, 1)$, and if $\xi = 0$ any point of the whole segment $(0, r)$, with $r \in [0, 1]$, is a fixed point. Moreover, the following holds:

- If $2\alpha\gamma > Na_iq_i$, $i = 0, 1$, then two more boundary-fixed points exist, namely $E_0^* = (x_0^*, 0)$ with x_0^* given by Equation (10) and $E_1^* = (x_1^*, 1)$ with x_1^* given by Equation (13).
- If $\xi < 0$, $a_0^2q_0 < a_1^2q_1$ and $k_0 < k < k_1$, where k_0 and k_1 are given by Equations (17) and (18), respectively, then an interior equilibrium $E^* = (x^*, r^*)$ exists with components given by Equation (16).

- If E_0^* exists (i.e. $2\alpha\gamma > Na_0q_0$) then for $\xi < 0$ E_0^0 is a saddle point with stable set along the invariant line $x = 0$ and unstable set along the invariant line $r = 0$.
- If E_1^* exists (i.e. $2\alpha\gamma > Na_1q_1$), then for $\xi < 0$ E_1^0 is an unstable node.
- E_0^* is a stable node if $k < k_0$ and $Na_0q_0/(2\gamma) < \alpha < Na_0q_0/(2\gamma) + 2$. At $k = k_0$, it undergoes a transcritical bifurcation at which it merges with E^* , at $\alpha < Na_0q_0/(2\gamma) + 2$ it undergoes a flip bifurcation along the invariant line $r = 0$.
- E_1^* is a stable node if $k > k_1$ and $Na_1q_1/(2\gamma) < \alpha < Na_1q_1/(2\gamma) + 2$. At $k = k_1$, it undergoes a transcritical bifurcation at which it merges with E^* , at $\alpha < Na_1q_1/(2\gamma) + 2$ it undergoes a flip bifurcation along the invariant line $r = 1$.
- The interior-fixed point E^* is stable if $k_0 < k < k_1$ and $\beta_F < \beta < \beta_{NS}$, where β_{NS} and β_F are given by Equations (30) and (32), respectively; hence, the range of stability is nonempty provided that $\alpha\gamma\xi > -k(a_1^2q_1 - a_0^2q_0)$.

It is worth highlighting that condition $\beta_F < \beta_{NS}$ is equivalent to $k < (\alpha x^*/4)$ that, being $k > k_0$, is verified for $1 + \alpha - Na_0q_0/(2\gamma) < 5$, that is true when the restriction Equation (9) to $r = 0$ has bounded dynamics, i.e. when $1 + \alpha - Na_0q_0/(2\gamma) \leq 4$. The stability range of the interior equilibrium E^* , as the parameter β varies, is shown by the bifurcation diagrams in Figure 2, where the supercritical flip and Neimark–Sacker bifurcations, through which the equilibrium loses its stability for decreasing and increasing values of β respectively, can be clearly seen.




Figure 2. Left column: bifurcation diagrams for $\beta \in [0, 350]$ showing the asymptotic dynamics of the two state variables x and r . parameters: $\alpha = 2.7$, $y = 1$, $N = 15$, $a_0 = 1.05487$, $a_1 = 0.5$, $q_0 = 0.01$, $q_1 = 0.2$, $k = 6.7$ and $\xi = -0.056$. Right column: attractors on the (x, r) phase space for different values of β . The other parameters are as in the left column.

4. Global dynamics

In this section, we propose further analytical results and numerical explorations of some global dynamic scenarios of the discrete dynamical system Equation (7) under the constraints on the parameters imposed by the economic and ecologic meaning of the model. The numerical simulations will confirm the analytical results on local stability and bifurcations given in the previous section and will give some snapshots about global dynamic behaviours.

Let us start by commenting the bifurcation diagrams presented in Figure 2. They show the long run dynamics of the model varying the intensity of choice parameter β , that represents the evolutionary propensity to switch to the more profitable technology. In evolutionary models a common occurrence is that an increase of the value of β leads to instability and complex dynamics, see e.g. [14]. However, this is not necessarily true in the case of the evolutionary model studied herein. Indeed, starting from values close to zero, as β increases we observe a transition from oscillatory dynamics (periodic or chaotic) towards non-oscillatory dynamics through a cascade of period-halving bifurcations leading to the stability of the fixed point E^* . Moreover, we have a range of the values of β , namely $\beta \in (\beta_F, \beta_{NS})$ as stated in Proposition 1, such that the fixed point E^* is stable. Furthermore, for $\beta > \beta_{NS}$, E^* becomes unstable again through a supercritical Neimark-Sacker bifurcation, at which a stable invariant closed curve is created around E^* , whose amplitude increases as β increases, see Figure 2 right panel. So, this example underlines the unusual result that low values of intensity of choice, as well as high values, lead to instability of the fixed point E^* with the creation of periodic or quasi-periodic or even chaotic attractors, whereas intermediate levels of the intensity of choice are required for the local asymptotic stability of E^* .

Other interesting results regard the evolutionary dominance of one of the two harvesting strategies. On the basis of the analytical results in the previous section, we know that as long as E^* exists, then the two fixed points E_0^* and E_1^* are transversally unstable. At first sight, this may suggest that an evolutionary dominant strategy, or equivalently an attractor along the invariant lines $r = 0$ or $r = 1$, exits if and only if the interior fixed point E^* is unfeasible and at the same time an attractor with $r \in (0, 1)$ exits if and only if the interior fixed point E^* is feasible. Instead, interior attractors (cyclic or chaotic) may exist even for $k > k_1$, i.e. after the transcritical bifurcation at which the equilibrium E^* merges with E_1^* and becomes unfeasible. Moreover, convergence towards the invariant line $r = 0$ may occur even when the equilibrium E^* is feasible, i.e. $k_0 < k < k_1$. These two occurrences are stated by the following two Propositions, respectively.

Proposition 2. Consider map Equation (7). Let $\xi < 0$ and the other parameters are positive and such that E_1^ exists and it is unstable along the manifold $r = 1$, and a period-2 cycle, say $\mathcal{C}_1^2 = \{(x_1^*, 1), (x_1^{*2}, 1)\}$, exists on the invariant set $r = 1$, as the result of the period-doubling bifurcation of E_1^* . If the following condition holds*

$$\xi - \frac{k(a_0^2 q_0 - a_1^2 q_1)}{2\alpha\gamma} < \frac{k(a_0^2 q_0 - a_1^2 q_1)(2\gamma\alpha - Na_1 q_1)}{8\alpha\gamma^2} - \xi < 0 \quad (33)$$

then the fixed point E_1^ has stable transverse invariant manifold and period-2 cycle \mathcal{C}_1^2 is transversely unstable. The contrary cannot occur.*

PROOF. See Appendix A. □

Proposition 2 hints at an interesting dynamic scenarios, confirmed by the numerical simulations shown in Figure 3. In this case, although there are no interior-fixed points and the border equilibrium E_1^* is transversely stable, the dynamics of the model can still converge in the long run to a stable inner attractor, a stable period-2 cycle C_1^2 shown in Figure 3 (left panel). This evidence indicates that the stability of the transverse invariant manifold of E_1^* does not imply the predominance by evolutionary pressure of the standard (or intensive) technology.

The situation is different (and in some sense reverted) when we consider the invariant line $r = 0$, as stated in the following proposition.

Proposition 3. Consider map Equation (7). Let $\xi < 0$ and the other parameters are positive and such that E_0^ exists and it is unstable along the invariant manifold $r = 0$, and a period-2 cycle, say $C_0^2 = \{(x_0^{*1}, 0), (x_0^{*2}, 0)\}$, exists on the invariant set $r = 0$, as the result of the period-doubling bifurcation of E_0^* . If the following condition holds*

$$\xi - \frac{k(a_0^2q_0 - a_1^2q_1)}{2\alpha\gamma} < \frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - \xi < 0 \tag{34}$$

then the fixed point E_0^ is transversely unstable and period-2 cycle C_0^2 has a stable transverse invariant manifold. The contrary cannot occur.*

PROOF. See Appendix B. □

The proposition 3 provides an interesting result highlighted by the numerical simulations shown in Figure 3 (right panel), where, although the border equilibrium E_0^* is




Figure 3. Left panel: the grey region is the basin of attraction of the inner period-2 cycle C_1^2 , while the fixed point E_1^* is transversely attractive. The point E^* is unfeasible, i.e. it lies in the region $r > 1$. Parameters' values: $\alpha = 3.1, k = 1, N = 8, q_1 = 1, q_0 = 0.01, \gamma = 4, a_0 = 1.1, a_1 = 1, \beta = 10, \xi = -0.0415$. Right panel: the grey region is the basin of attraction of the transversely stable fixed point E^* and the white region is the basin of attraction of the period-2 cycle C_0^2 . Parameters' values: $\alpha = 2.6, k = 1, N = 8, q_1 = 1, q_0 = 0.1, \gamma = 2, a_0 = 1.1, a_1 = 1, \beta = 80.615, \xi = -0.095$.

transversally unstable, the 2-cycle C_0^2 is locally asymptotically stable and coexists with the locally asymptotically stable interior fixed point E^* .

Numerical investigations suggest that this dynamic scenario occurs due to a specific sequence of bifurcations. In particular, a 2-cycle in the region with negative r , let us name it C^2 , undergoes a transcritical bifurcation, merging with the 2-cycle C_0^2 originated by a period-doubling bifurcation of the fixed point E_0^* , and becomes feasible. After the bifurcation, C_0^2 becomes local asymptotically stable, while C^2 is a saddle 2-cycle, and its one-dimensional stable manifold marks the boundary separating the basins of attraction of E^* and C_0^2 . Then, changing the values of the parameters in a suitable way, the 2-cycle C^2 disappears through a subcritical flip bifurcation at which E^* loses stability and becomes a saddle fixed point. These bifurcations occur before the merging of E^* with E_0^* . This underlines that agents can select the environmentally friendly technology even when E^* is a feasible fixed point and E_0^* is transversely unstable. These results point out a quite peculiar property of the considered evolutionary model. In fact, the instability of the inner fixed point of the model may lead to an increase in the propensity of the agent to adopt the environmentally friendly technology. These scenarios do not occur on the continuous (or hybrid) setting of the model analysed in [15].

The basins of attraction in Figure 3 underline further interesting properties of the dynamics of the model. In particular, from the right panel in Figure 3, it is possible to observe that the transverse unstable manifold of E_0^* belongs to the basin of attraction of E^* ; hence, such a basin has a contact with the invariant line $r = 0$ at the point E_0^* . This implies that all the pre-images of E_0^* along the invariant line $r = 0$, computed according to the restriction Equation (9), represent tongues at which the basin of E^* has a contact with line $r = 0$. In the figure only some of them are visible, but infinitely many exist and accumulate near E_0^* . The fine structure of these tongues is quite complicated and will be analysed in future works.

It is worth observing that the mechanisms that lead to the evolutionary-dominant environmentally friendly technology when the fixed point E^* is feasible, can be even different from the described one and, as shown in the following, can be due to the existence of non-topological Milnor attractors on the invariant line $r = 0$. For example, the bifurcation diagrams in Figure 4, obtained varying parameter a_0 in the range $[0.5, 1.3]$, show another case such that an attracting invariant set \mathcal{A}_s , laying on the axis $r = 0$, exists even when the interior equilibrium point E^* is feasible. Measure-theoretic arguments about the transverse attractiveness of the invariant set \mathcal{A}_s can be used to provide an explanation of this dynamic phenomenon. In particular, we make use of the transverse Lyapunov exponent (see e.g. [2], [8]) defined as:

$$\Lambda_{\perp} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^N \ln |\nu_{\perp}(x_n)|, \quad (35)$$

where $\{x_n = f_0^n(x_0), n \geq 0\}$ is a trajectory embedded in \mathcal{A}_s and $\nu_{\perp}(x_n)$ is the transverse eigenvalue computed in x_n . Precisely, when \mathcal{A}_s is a k -cycle C_0^k , its transverse attractiveness is measured by the product of the transverse eigenvalues $\nu_{\perp}^k = \prod_{i=1}^k \nu_{\perp}(x_i)$ and, if $\Lambda_{\perp}(C^k) = k^{-1} \ln |\nu_{\perp}^k| < 0$, then \mathcal{A}_s is a topological attractor. While, when the attractor \mathcal{A}_s is chaotic and so includes infinitely many cycles densely distributed within it, each one characterized by its own transversal Lyapunov exponent, transverse attractiveness of \mathcal{A}_s can be measured by the spectrum of the Lyapunov exponents. This is defined, see e.g. [8], as

$$\Lambda_{\perp}^{\min} \leq \dots \leq \Lambda_{\perp}^{\text{nat}} \leq \dots \leq \Lambda_{\perp}^{\max}, \quad (36)$$

where Λ_{\perp}^{\min} and Λ_{\perp}^{\max} are the Lyapunov exponents of the most attractive and the most repelling cycles in \mathcal{A}_s , respectively. Moreover, the *natural Lyapunov exponent* $\Lambda_{\perp}^{\text{nat}}$ is computed along a generic aperiodic trajectory embedded in \mathcal{A}_s , and it measures transversal attractiveness on *average*. In other words, $\Lambda_{\perp}^{\text{nat}}$ carries contributions to attractiveness from all the trajectories in \mathcal{A}_s , giving the mean local behaviour in its neighbourhood (see e.g. [19]). If \mathcal{A}_s contains at least one transversely repelling cycle with a dense set of pre-images embedded in \mathcal{A}_s , that is $\Lambda_{\perp}^{\max} > 0$, then there is no neighbourhood of \mathcal{A}_s containing only points whose ω -limit set belongs to \mathcal{A}_s . According to the theorem stated in [1], the latter inequality implies that the one-dimensional invariant chaotic set cannot be a Lyapunov attractor in the two-dimensional space because of the transversely unstable set of the period-2 cycle as well as its pre-images. If the inequality $\Lambda_{\perp}^{\text{nat}} < 0$ holds also, thus the set \mathcal{A}_s attracts a positive measure set of points that converges to it. It follows that this is an attractor in Milnor sense (see [18]). A large number of results about global attractiveness of invariant manifold of lower dimension than the total phase space can be found in the literature, see e.g. [1,2,8].

In the right panel in Figure 4, we present both the transverse Lyapunov exponent characterizing transverse attractiveness of \mathcal{A}_s and the transverse Lyapunov exponent of to the period-2 cycle C_0^2 . For suitable values of the aggregate parameter $\mu = 1 + \alpha - Na_0q_0/(2\gamma)$, at which the 2-cycle of the logistic map undergoes the homoclinic bifurcation due to which 2-cyclic chaotic intervals are obtained by the merging of 4-cyclic chaotic intervals, pure chaos exists in \mathcal{A}_s (see e.g [16]). For example, for $\alpha = 2.7$, $N = 20$, $\gamma = 1$, $q_0 = 0.01$ and $a_0 = a_c = 1.07428$, we detect the presence of a Milnor attractor when $\mu = \bar{\mu} \approx 3.5925721841$, value at which $\Lambda_{\perp}^{\text{nat}}(\bar{\mu}) < 0$, while $\Lambda_{\perp}^{\max} \geq \Lambda_{\perp}(C_0^2) > 0$. Note that the parameter β is the so-called normal parameter, i.e. it affects only the transverse stability of \mathcal{A}_s and does not have influence on the dynamics inside the invariant set \mathcal{A}_s . Setting $\mu = 1 + \alpha - Na_0q_0/(2\gamma) = \bar{\mu}$ and varying β , we vary the spectrum of Lyapunov exponents, i.e. we change the topological property of the invariant set \mathcal{A}_s which turns to be a chaotic saddle, a non-topological Milnor attractor and a topological Lyapunov stable set.




Figure 4. Bifurcation diagram of x vs. a_0 (left panel) and r vs. a_0 (centre panel) varying $a_0 \in (a_1, a_1\sqrt{q_1/q_0})$. Dashed lines represent paths of both x^* and r^* . We can notice that the asymptotic dynamics is enclosed along the invariant axis $r = 0$ even for $r^*(a_0) > 0$. This is due to the transverse attractiveness of some subsets of the invariant axes $r = 0$, while both the fixed point E_0^* and the period-2 cycle C_0^2 are transversally repelling. For $\mu = 1 + \alpha - Na_0q_0/(2\gamma) \approx 3.5925721841$ such that $a_0 = a_c = 1.07428$, the attractor \mathcal{A}_s of the logistic map on $r = 0$ is characterized by pure chaos. Since $\Lambda_{\perp}^{\text{nat}}(\bar{\mu}) \approx -0.0034703 < 0$ while $\Lambda_{\perp}(C_0^2) > 0$, it results that \mathcal{A}_s is a non-topological Milnor attractor. Parameters' values are as in Figure 2, but $N = 20$ and $\beta = 5$.




Figure 5. Upper panel: a trajectory in the phase space. Lower panel: a time series of $r(t)$. Parameters' values are as in Figure 2, but $N = 20$ and $\beta = 5$.

To sum up, we can say that \mathcal{A}_s can attract a set of positive Lebesgue measures even when it contains repelling cycles together with dense sets of their pre-images.

In the last part of the section, we point out that numerical simulations of the model can show dynamic scenarios that are difficult to infer analytically. For example, in Figure 5 we observe a chaotic attractor in the region $r \in (0, 1)$ where both harvesting strategies coexist, although there is a prevalence of the environmentally friendly one. The time series $r(t)$ shows an apparently stochastic behaviour, which typically characterizes the evolutionary dynamics driven by a replicator equation.

To conclude this section, we want to emphasize that the conducted investigation underlines that the class of evolutionary games, represented by the discrete-time model considered here, reveals some interesting and economically insightful dynamics, such as non-topological Milnor attractors, that are not observable in other and more simple evolutionary games, see e.g. [14].

5. Conclusions

In this paper, we have analysed some local and global dynamical properties of a map that represents the discrete-time version of an evolutionary game model proposed in [15] to describe a fishery where a pool of fishermen can select between two different technologies of harvesting, that is an environmentally friendly technology and a standard technology. The analysis of the dynamics underlines interesting scenarios that cannot occur on the continuous version of the model. In particular, Neimark–Sacker bifurcations and cascade of period-doubling bifurcations can lead to quite complicated, even chaotic, dynamics. Moreover, a deeper analytical and numerical analysis reveals the existence of Milnor attractors. Those attractors have interesting economic implications not only for the model proposed here, but also for an entire class of evolutionary games that this model could represent. For this reason, the analysis and the presence of the Milnor attractors and related

study of riddled basins for this class of evolutionary games represent an interesting aspect that deserve further investigation.

Acknowledgements

We thank a Guest Editor and an anonymous Referee for useful comments and suggestions. This work has been performed within the framework of COST Action IS1104 ‘The EU in the new economic complex geography: models, tools and policy evaluation’ and under the auspices of GNFM, Gruppo Nazionale di Fisica Matematica (Italy).

Disclosure statement

No potential conflict of interest was reported by the authors.

Notes

1. Email: gian.bischi@uniurb.it
2. Email: d.radi@univpm.it
3. For sake of completeness, we stress that if $\xi > 0$, i.e. $c_0 > c_1$, then $x^* > 0$ provided that $a_0^2 q_0 > a_1^2 q_1$, that together with the condition $a_0 q_0 < a_1 q_1$ gives $(a_1^2 q_1 / a_0) < a_0 q_0 < a_1 q_1$, which is a nonempty set being $a_0 > a_1$.

References

- [1] J.C. Alexander, J.A. Yorke, Z. You, and I. Kan, *Riddled basins*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. 2(4) (1992), pp. 795–813. doi:10.1142/S0218127492000446.
- [2] P. Ashwin, J. Buescu, and I. Stewart, *From attractor to chaotic saddle: A tale of transverse instability*, Nonlinearity 9(3) (1996), pp. 703–737. doi:10.1088/0951-7715/9/3/006.
- [3] G.I. Bischi, F. Lamantia, and D. Radi, *A prey–predator fishery model with endogenous switching of harvesting strategy*, Appl. Math. Comput. 219(20) (2013), pp. 10123–10142. doi:10.1016/j.amc.2013.02.061.
- [4] G.I. Bischi, F. Lamantia, and D. Radi, *Multi-species exploitation with evolutionary switching of harvesting strategies*, Nat. Resour. Model. 26(4) (2013), pp. 546–571. doi:10.1111/nrm.12013.
- [5] G.I. Bischi, F. Lamantia, and L. Sbragia, *Competition and cooperation in natural resources exploitation: An evolutionary game approach*, in *Game Practice and the Environment*, C. Carraro and V. Fragnelli, eds., Edward Elgar Publishing, Cheltenham, UK, 2009, pp. 187–211.
- [6] G.I. Bischi, F. Lamantia, and F. Tramontana, *Sliding and oscillations in fisheries with on-off harvesting and different switching times*, Commun. Nonlinear Sci. Numer. Simul. 19(1) (2014), pp. 216–229.
- [7] F. Brauer, and C. Castillo-Chavez, *Mathematical Models in Population Biology and Epidemiology*, Springer-Verlag, New York, 2001, volume Texts in Applied Mathematics 40.
- [8] J. Buescu, *Exotic Attractors*, Birkh’user, Basel, 1997.
- [9] A. Cabrales, and J. Sobel, *On the limit points of discrete selection dynamics*, J. Econom. Theory 57(2) (1992), pp. 407–419. doi:10.1016/0022-0531(92)90043-H.
- [10] J.M. Cushing, *Some discrete competition models and the competitive exclusion principle*, J. Difference Equ. Appl. 10(13–15) (2004), pp. 1139–1151. doi:10.1080/10236190410001652739.
- [11] S.N. Elaydi, *An introduction to difference equations*, Springer, New York, 1995.
- [12] S.A.H. Geritz, and E. Kisdi, *On the mechanistic underpinning of discrete-time population models with complex dynamics*, J. Theoret. Biol. 228(2) (2004), pp. 261–269. doi:10.1016/j.jtbi.2004.01.003.
- [13] W.M. Getz, and R.G. Haight, *Population Harvesting: Demographic Models of Fish, Forest, and Animal Resources*, Princeton University Press, Princeton, NJ, 1989, 391 pp.
- [14] J. Hofbauer, and K. Sigmund, *Evolutionary game dynamics*, Bull. (N.S.) Amer. Math. Soc. 40(4) (2003), pp. 479–520. doi:10.1090/S0273-0979-03-00988-1.
- [15] F. Lamantia, and D. Radi, *Exploitation of renewable resources with differentiated technologies: an evolutionary analysis*, Math. Comput. Simulation 108 (2015), pp. 155–174. doi:10.1016/j.matcom.2013.09.013.

- [16] Y.L. Maistrenko, V.L. Maistrenko, A. Popovich, and E. Mosekilde, *Transverse instability and riddled basins in a system of two coupled logistic maps*, Phys. Rev. E 57(3) (1998), pp. 2713–2724. doi:10.1103/PhysRevE.57.2713.
- [17] R.M. May, *Biological populations obeying difference equations: stable points, stable cycles, and chaos*, J. Theoret. Biol. 51(2) (1975), pp. 511–524. doi:10.1016/0022-5193(75)90078-8.
- [18] J. Milnor, *On the concept of attractor: Correction and remarks*, Comm. Math. Phys. 102(3) (1985), pp. 517–519. doi:10.1007/BF01209298.
- [19] Y. Nagai, and Y.-C. Lai, *Periodic-orbit theory of the blowout bifurcation*, Phys. Rev. E 56(4) (1997), pp. 4031–4041. doi:10.1103/PhysRevE.56.4031.
- [20] P. Taylor, and L. Jonker, *Evolutionary stable strategies and game dynamics*, Math. Biosci. 40(1–2) (1978), pp. 145–156. doi:10.1016/0025-5564(78)90077-9.

Appendix A

Let us rewrite the map T as follows:

$$T : \begin{cases} x' = Ax - Bx^2 + Cxr \\ r' = \frac{r}{r+(1-r)e^{Dx-\beta\xi}} \end{cases}, \quad (37)$$

where

$$A = 1 + \alpha - \frac{Na_0q_0}{2\gamma}; \quad B = \frac{\alpha}{k}; \quad C = \frac{N}{2\gamma}(a_0q_0 - a_1q_1); \quad D = \beta \frac{a_0^2q_0 - a_1^2q_1}{4\gamma} \quad (38)$$

and let us consider the second iterate of the map T , i.e.

$$T^2 : \begin{cases} x' = (Ax - Bx^2 + Cxr) \left(A + C \frac{r}{r+(1-r)e^{Dx-\beta\xi}} \right) - B(Ax - Bx^2 + Cxr)^2 \\ r' = \frac{\frac{r}{r+(1-r)e^{Dx-\beta\xi}}}{\frac{r}{r+(1-r)e^{Dx-\beta\xi}} + \left(1 - \frac{r}{r+(1-r)e^{Dx-\beta\xi}} \right) e^{D(Ax - Bx^2 + Cxr) - \beta\xi}} \end{cases} \quad (39)$$

Its restriction to the invariant line $r = 1$ is

$$T^2|_{r=1} : \begin{cases} x' = F(Fx - Bx^2) - B(Fx - Bx^2)^2 \\ r' = 1 \end{cases}, \quad (40)$$

where

$$F = A + C = 1 + \alpha - \frac{Na_1q_1}{2\gamma}. \quad (41)$$

The map T^2 can have at most four fixed points, given by the solutions of the equation

$$x(B^3x^3 - 2FB^2x^2 + FB(1+F)x + 1 - F^2) = 0 \quad (42)$$

from which we obtain

$$E_1^0 = (0, 1), E_1^* = (x_1^*, 1), E_1^{*1} = (x_1^{*1}, 1) \text{ and } E_1^{*2} = (x_1^{*2}, 1), \quad (43)$$

where x_1^* is given by Equation (13) and

$$x_1^{*1} = \frac{1 + F + \sqrt{F^2 - 3 - 2F}}{2B}, \quad x_1^{*2} = \frac{1 + F - \sqrt{F^2 - 3 - 2F}}{2B}.$$

Assuming the existence of period-2 cycle $\{(x_1^{*1}, 1), (x_1^{*2}, 1)\}$ of T is equivalent to the existence of E_1^{*1} and E_1^{*2} for T^2 which, requiring also $x_1^{*1} > 0, x_1^{*2} > 0$, implies $F > 3$, i.e. $\alpha - 2 > Na_1q_1/(2\gamma)$. The Jacobian matrix associated to T^2 along the restriction $r = 1$ is

$$J^2(x, 1) = \begin{bmatrix} (F - 2Bx)(F - 2BFx + 2B^2x^2) & J_{12}^2(x, 1) \\ 0 & J_{22}^2(x, 1) \end{bmatrix}, \tag{44}$$

where $J_{22}^2(x, 1) = e^{D(Fx - Bx^2 + x) - 2\beta\xi}$, from which we have the condition for transverse stability of E_1^{*1} and E_1^{*2} , given by $J_{22}^2(x_1^{*1}, 1) = J_{22}^2(x_1^{*2}, 1) < 1$. By trivial algebra, we obtain the condition

$$D\left(\frac{F + 1}{B}\right) - 2\beta\xi < 0. \tag{45}$$

Substituting for D, F and B , we obtain

$$\frac{k(a_0^2q_0 - a_1^2q_1)(4\gamma + 2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} - 2\xi < 0. \tag{46}$$

By similar calculation, the condition to have stable the transverse manifold of the fixed point E_1^* of T^2 is given by

$$\frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} - \xi < 0. \tag{47}$$

From conditions Equations (46) and (47), it follows that the transverse invariant manifold of E_1^* is stable and the transverse invariant manifolds of E_1^{*1} and E_1^{*2} are unstable if and only if

$$\frac{k(a_0^2q_0 - a_1^2q_1)(4\gamma + 2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} - 2\xi > 0 > \frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} - \xi \tag{48}$$

which can be rewritten as follows:

$$\xi - \frac{k(a_0^2q_0 - a_1^2q_1)}{2\alpha\gamma} < \frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} - \xi < 0. \tag{49}$$

By simple considerations, it is easy to note that conditions Equation (49) and condition $\alpha - Na_1q_1/(2\gamma) > 2$ required for the existence of E_1^{*1} and E_1^{*2} identify a nonempty set of the parameter space.

Moreover, from stability condition Equations (46) and (47), it is easy to note that in order to have the instability of the transverse invariant manifold of E_1^* and the stability of

the transverse invariant manifolds of E_1^{*1} and E_1^{*2} is required

$$\frac{k(a_0^2 q_0 - a_1^2 q_1)(4\gamma + 2\gamma\alpha - Na_1 q_1)}{8\alpha\gamma^2} - 2\xi < 0 < \frac{k(a_0^2 q_0 - a_1^2 q_1)(2\gamma\alpha - Na_1 q_1)}{8\alpha\gamma^2} - \xi \quad (50)$$

which can be rewritten as:

$$\frac{k(a_0^2 q_0 - a_1^2 q_1)}{2\alpha\gamma} - \xi < \xi - \frac{k(a_0^2 q_0 - a_1^2 q_1)(2\gamma\alpha - Na_1 q_1)}{8\alpha\gamma^2} < 0. \quad (51)$$

Since throughout this paper, we always assume $a_0^2 q_0 - a_1^2 q_1 < 0$ and $\alpha - Na_1 q_1 / (2\gamma) > 2$ is required for the existence of E_1^{*1} and E_1^{*2} , condition Equation (51) implies that $(|\cdot|)$ is the absolute value of \cdot)

$$|\xi| > \left| \frac{k(a_0^2 q_0 - a_1^2 q_1)(2\gamma\alpha - Na_1 q_1)}{8\alpha\gamma^2} \right| > \left| \frac{k(a_0^2 q_0 - a_1^2 q_1)}{2\alpha\gamma} \right| > |\xi| \quad (52)$$

which is a contradiction. It follows that condition Equation (51) cannot be satisfied.

Since the condition to have stable (or unstable) transversally manifold of the period 2-cycle $\{(x_1^{*1}, 0), (x_1^{*2}, 0)\}$ of the map T is equivalent to condition to have stable transverse invariant manifold of each of the two fixed points E_1^{*1} and E_1^{*2} of T^2 and the condition to have stable (or unstable) transverse invariant manifold of the fixed point E_1^* are the same for T and T^2 , the claim of the proposition follows.

Appendix B

Let us consider the restriction of T^2 , defined in Appendix A (see Equation (37)), on the invariant line $r = 0$

$$T^2|_{r=0} : \begin{cases} x' = A(Ax - Bx^2) - B(Ax - Bx^2)^2 \\ r' = 0 \end{cases} \quad (53)$$

It has at most four fixed points given by the solutions of the equation

$$x(B^3 x^3 - 2B^2 A x^2 + AB(1+A)x + 1 - A^2) = 0 \quad (54)$$

from which we obtain:

$$E_0^0 = (0, 0), E_0^* = (x_0^*, 0), E_0^{*1} = (x_0^{*1}, 0) \text{ and } E_0^{*2} = (x_0^{*2}, 0), \quad (55)$$

where x_0^* is given in Equation (10), and

$$x_0^{*1} = \frac{1 + A + \sqrt{A^2 - 3 - 2A}}{2B}, \quad x_0^{*2} = \frac{1 + A - \sqrt{A^2 - 3 - 2A}}{2B}.$$

Assuming the existence of period-2 cycle $\{(x_0^{*1}, 0), (x_0^{*2}, 0)\}$ of T is equivalent to the existence of E_0^{*1} and E_0^{*2} for T^2 which, requiring also $x_0^{*1} > 0$ and $x_0^{*2} > 0$, implies $A > 3$,

i.e. $\alpha - 2 > Na_0q_0/(2\gamma)$. The Jacobian matrix of T^2 along $r = 0$ is

$$J^2(x, 0) = \begin{bmatrix} (A - 2Bx)(A - 2BAx + 2B^2x^2) & J_{12}^2(x, 0) \\ 0 & J_{22}^2(x, 0) \end{bmatrix}, \tag{56}$$

where $J_{22}^2(x, 0) = e^{-D(Ax - Bx^2 + x) + 2\beta\xi}$, from which the condition for transverse stability of E_0^{*1} and E_0^{*2} is $J_{22}^2(x_0^{*1}, 0) = J_{22}^2(x_0^{*2}, 0) < 1$. By trivial algebra, we obtain the condition

$$D\left(\frac{1+A}{B}\right) - 2\beta\xi > 0. \tag{57}$$

Substituting for D , A and B , we obtain

$$\frac{k(a_0^2q_0 - a_1^2q_1)(4\gamma + 2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - 2\xi > 0. \tag{58}$$

By similar calculation, the condition for transverse stability of the fixed point E_0^* of T^2 is given by

$$\frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - \xi > 0. \tag{59}$$

From the conditions [Equations \(58\) and \(59\)](#), it follows that transverse invariant manifold of E_0^* is unstable and transverse invariant manifolds of E_0^{*1} and E_0^{*2} are stable if and only if

$$\frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - \xi < 0 < \frac{k(a_0^2q_0 - a_1^2q_1)(4\gamma + 2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - 2\xi \tag{60}$$

which can be rewritten as follows:

$$\xi - \frac{k(a_0^2q_0 - a_1^2q_1)}{2\alpha\gamma} < \frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - \xi < 0. \tag{61}$$

By simple considerations, it is easy to note that conditions [Equation \(61\)](#) and condition $\alpha - Na_0q_0/(2\gamma) > 2$ required for the existence of E_0^{*1} and E_0^{*2} identify a nonempty set of the parameter space.

Moreover, from stability condition [Equations \(58\) and \(59\)](#), it is easy to note that in order to have the stability of the transverse invariant manifold of E_0^* and the instability of the transverse invariant manifolds of E_0^{*1} and E_0^{*2} is required

$$\frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - \xi > 0 > \frac{k(a_0^2q_0 - a_1^2q_1)(4\gamma + 2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} - 2\xi \tag{62}$$

which can be rewritten as follows:

$$0 > -\frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_0q_0)}{8\alpha\gamma^2} + \xi > \frac{k(a_0^2q_0 - a_1^2q_1)}{2\alpha\gamma} - \xi. \quad (63)$$

Since throughout this paper, we always assume $a_0^2q_0 - a_1^2q_1 < 0$ and $\alpha - Na_1q_1/(2\gamma) > 2$ is required for the existence of E_1^{*1} and E_1^{*2} , condition Equation (51) implies that $(|\cdot|$ is the absolute value of \cdot)

$$|\xi| > \left| \frac{k(a_0^2q_0 - a_1^2q_1)(2\gamma\alpha - Na_1q_1)}{8\alpha\gamma^2} \right| > \left| \frac{k(a_0^2q_0 - a_1^2q_1)}{2\alpha\gamma} \right| > |\xi| \quad (64)$$

which is a contradiction. It follows that condition Equation (63) cannot be satisfied.

Since the condition to have stable (or unstable) transversally manifold of the period 2-cycle $\{(x_1^{*1}, 0), (x_1^{*2}, 0)\}$ of the map T is equivalent to condition to have stable transverse invariant manifold of each of the two fixed points E_1^{*1} and E_1^{*2} of T^2 and the condition to have stable (or unstable) transverse invariant manifold of the fixed point E_1^* are the same for T and T^2 , the claim of the proposition follows.